#### **BIO-COX - salinomycin** granule

Alpharma Inc. Animal Health

Brand of salinomycin sodium
Type A Medicated Article
For use in broiler, roaster, replacement
(breeder and layer) chickens, and quail only

## **ACTIVE DRUG INGREDIENT**

Salinomycin equivalent to 60 grams salinomycin sodium activity per pound

For use in manufacturing medicated feeds only

## Broiler, roaster and replacement (breeder and layer) chickens

Use Bio-Cox 60 to provide salinomycin sodium concentrations ranging from 40-60 g/ton (0.0044% to 0.0066%) for the prevention of coccidiosis in broiler, roaster and replacement chickens caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions.

## **QUAIL**

Use Bio-Cox 60 to provide salinomycin sodium concentrations of 50 g/ton (0.0055%) for the prevention of coccidiosis in quail caused by *Eimeria dispersa* and *E. lettyae*.

#### MIXING DIRECTIONS

Thoroughly mix Bio-Cox 60 with non-medicated feed to provide the level of salinomycin sodium per ton as indicated in the table below.

| Salinomycin Sodium Activity | Amount of Bio-Cox 60 Type A Medicated<br>Article (60 g/lb) Per Ton of Feed |       |
|-----------------------------|--|-------|
| (grams per ton)             | lb   | grams |
| 40                          | 0.67   | 304   |
| 45                          | 0.75   | 341   |
| 50                          | 0.83   | 377   |
| 55                          | 0.92   | 418   |
| 60                          | 1.00   | 454   |

#### FEEDING DIRECTIONS

Feed continuously as the sole ration

#### **IMPORTANT**

Must be thoroughly mixed in feeds before use. Not approved for use with pellet binders.

# **CAUTION**

May be fatal if fed to adult turkeys or to horses.

# WARNING

Do not feed to laying hens producing eggs for human consumption. When mixing and handling Bio-Cox 60 premix, use protective clothing, impervious gloves, eye protection, and an approved dust respirator. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Store in a cool, dry place

NADA #128-686, Approved by FDA

Marketed by Alpharma Inc. Bridgewater, New Jersey 08807 **Net wt 50 LB (22.68 kg)** 



Trademarks registered by Alpharma Inc.

Made in USA 400321 0907

# PRINCIPAL DISPLAY PANEL - 50 LB LABEL

Bio-Cox®
60
ALPHARMA
Brand of salinomycin sodium
Type A Medicated Article
Important: See back panel for use directions
Net wt 50 LB (22.68 kg)
ALPHARMA®

Bio-Cox<sup>®</sup>



